

Determination of structure of glycans and glycan interactions with proteins by NMR

Antonio Molinaro




molinaro@unina.it


*Department of Chemical Sciences, University of Napoli "Federico II"
Department of Chemistry, School of Science, Osaka University*

Why glycans are puzzling, why are they not obvious?


RNA




DNA






Protein



Glycan



A Open Form**B** Pyranose (5→1)**C** Furanose (4→1)

Open, pyranose and furanose forms of an aldose, showing the many equivalent OH groups. The OH groups can be modified by $-\text{PO}_4^{-3}$, $-\text{SO}_4^{-2}$, $-\text{NH}_2$, $-\text{CO}-\text{CH}_3$, $-\text{NH}-\text{CO}-\text{CH}_3$, $-\text{H}$.

Why glycans are puzzling, why are they not obvious?

Structure determination of a glycan chain: wet organic chemistry for every step



Available online at www.sciencedirect.com

ScienceDirect

Current Opinion in
Structural Biology



Solving the structural puzzle of bacterial glycome

Roberta Marchetti, Rosa Ester Forgione, Ferran Nieto Fabregat,
Cristina Di Carluccio, Antonio Molinaro and Alba Silipo



Contents lists available at ScienceDirect


Carbohydrate Polymers

journal homepage: www.elsevier.com/locate/carbpol

Liquid-state NMR spectroscopy for complex carbohydrate structural analysis: A hitchhiker's guide

Immacolata Speciale^a, Anna Notaro^a, Pilar Garcia-Vello^b, Flaviana Di Lorenzo^a,
Samantha Armiento^b, Antonio Molinaro^b, Roberta Marchetti^b, Alba Silipo^b,
Cristina De Castro^{a,c}

- ❖ Quali-quantitative analysis (**OC**, GC-MS, NMR)
- ❖ Absolute configuration (**OC**, GC-MS, NMR)
- ❖ Size of the ring (**OC**, GC-MS, NMR)
- ❖ Anomeric configuration (**OC**, NMR)
- ❖ Linkage analysis (**OC**, GC-MS, NMR)
- ❖ Monosaccharides sequence (**OC**, MALDI-MS, 2D NMR)
- ❖ Determination of non-carbohydrate appendages (**OC**, GC-MS, MALDI-MS, 2D NMR)



Typical Applications of NMR:

- **Structural (chemical) elucidation**

Natural product chemistry

Organic chemistry: Analytical tool of choice for synthetic chemists.

- **Study of dynamic processes**

Reaction kinetics.

Study of equilibrium (chemical or structural).

- **Structural (three-dimensional) studies**

Proteins

DNA/RNA

Polysaccharides

Drug design

- **Structure Activity Relationships (SAR) by NMR**

- **Medicine - Magnetic Resonance Imaging (MRI)**


NMR active nuclei

Group	I	II	IIIa	IVa	Va	VIa	VIIa	VIIIa	VIIIb	VIIIc	IB	IIB	III	IV	V	VI	VII	VIII
Period																		
1	1 H																	2 He
2	3 Li	4 Be											5 B	6 C	7 N	8 O	9 F	10 Ne
3	11 Na	12 Mg											13 Al	14 Si	15 P	16 S	17 Cl	18 Ar
4	19 K	20 Ca	21 Sc	22 Ti	23 V	24 Cr	25 Mn	26 Fe	27 Co	28 Ni	29 Cu	30 Zn	31 Ga	32 Ge	33 As	34 Se	35 Br	36 Kr
5	37 Rb	38 Sr	39 Y	40 Zr	41 Nb	42 Mo	43 Tc	44 Ru	45 Rh	46 Pd	47 Ag	48 Cd	49 In	50 Sn	51 Sb	52 Te	53 I	54 Xe
6	55 Cs	56 Ba	* 71 Lu	72 Hf	73 Ta	74 W	75 Re	76 Os	77 Ir	78 Pt	79 Au	80 Hg	81 Tl	82 Pb	83 Bi	84 Po	85 At	86 Rn
7	87 Fr	88 Ra	** 103 Lr	104 Unq	105 Unp	106 Unh	107 Uns	108 Uno	109 Mt	110 Uun	111 Uuu	112 Uub	113 Uut	114 Uuq	115 Uup	116 Uuh	117 Uus	118 Uuo
*Lanthanides	*		57 La	58 Ce	59 Pr	60 Nd	61 Pm	62 Sm	63 Eu	64 Gd	65 Tb	66 Dy	67 Ho	68 Er	69 Tm	70 Yb		
**Actinides	**		89 Ac	90 Th	91 Pa	92 U	93 Np	94 Pu	95 Am	96 Cm	97 Bk	98 Cf	99 Es	100 Fm	101 Md	102 No		
			Nuclear Spins															
			1/2	1	3/2	5/2	7/2	9/2										


Nucleus (odd atomic number): ^1H , ^{13}C , ^{15}N , ^{19}F , ^{31}P

How does NMR work?

A spinning charge creates a magnetic moment, so these nuclei can be thought of as tiny magnets.




In presence of a magnetic field
Magnetic moments precess and orient with or against the field




many nuclei

Line up with magnetic field




tip using RF pulse

Flip over




nuclei precess in xy plane
~few seconds



NMR Signal

Free Induction Decay
(FID)




Time

FT



NMR Signal



Frequency (Hz or MHz)




$$\nu_1 = \gamma B_0 / 2\pi.$$

¹³C NMR spectrum

$$\nu_1 = \gamma B_0 / 2\pi.$$

Chemical shift (ppm)



$$\delta \text{ (ppm)} = \frac{\nu_{\text{PEAK}} - \nu_{\text{REF}} \text{ (Hz)}}{\text{Freq of the nuclei (MHz)}} = \text{ppm}$$


- Depending on the *chemical environment* we have variations on the magnetic field that the nuclei feel, even for the same type of nuclei. It affects the local magnetic field.

$^1\text{H} - ^1\text{H}$ Coupling

Signals do not appear as single lines, sometimes they appear as multiple lines. This is due to $^1\text{H} - ^1\text{H}$ coupling (also called spin-spin splitting **or J-coupling**).


For this line, H_B is lined up **with** the magnetic field (adds to the overall magnetic field, so the line comes at higher frequency)

For this line, H_B is lined up **against** the magnetic field (subtracts from the overall magnetic field, so the line comes at lower frequency)




H_A is split into two lines because it feels the magnetic field of H_B .

H_B is split into two lines because it feels the magnetic field of H_A .




A and B are scalarly coupled nuclei

Doublet: 1 Adjacent Proton



Triplet: 2 Adjacent Protons



The $N + 1$ Rule


If a signal is split by N equivalent protons, it is split into $N + 1$ peaks.

n=0	1
n=1	1 1
n=2	1 2 1
n=3	1 3 3 1
n=4	1 4 6 4 1
n=5	1 5 10 10 5 1
n=6	1 6 15 20 15 6 1


Intensities




triplet n = 2




quartet n = 3



pentet n = 4



sextet n = 5



nonet n = 8


n	2^n	Multiplet Intensities
0	1	1
1	2	1 1
2	4	1 2 1
3	8	1 3 3 1
4	16	Pentet 1 4 6 4 1
5	32	Sextet 1 5 10 10 5 1
6	64	Septet 1 6 15 20 15 6 1
7	128	Octet 1 7 21 35 35 21 7 1
8	256	Nonet 1 8 28 56 70 56 28 8 1

^1H NMR

- ☞ Number of signals (number of non-equivalent H)
- ☞ Chemical shift (differences in chemical environment)
- ☞ Splitting or Coupling (number of neighboring H)
- ☞ Integration (relative number of H at each signal)

- Chemical shift data - tells us what kinds of protons we have.
- Integrals - tells us the ratio of each kind of proton in our sample.
- ^1H - ^1H coupling - tells us about protons that are near other protons.

- Number of signals
- Chemical shift
- Splitting or coupling
- Integration




^{13}C NMR

- ☞ Chemical shift is normally 0 to 220 ppm
- ☞ Similar factors affect the chemical shifts in ^{13}C as seen for H NMR
- ☞ Long relaxation times (excited state to ground state) mean no integration
- ☞ Number of peaks indicates the number of types of C



Ethyl ethanoate has 4 types of carbon so 4 peaks.
The most obvious peaks are the ester carbonyl at 190ppm
and the deshielded $-\text{CH}_2-$ attached to O at about 70 ppm.



Types of NMR experiments

● 1D Experiments:

- ^1H
- ^{13}C
- ^{31}P

● 2D Experiments:

- COSY (Correlation Spectroscopy)
- TOCSY (Total Correlation Spectroscopy)
- NOESY (Nuclear Overhauser effect spectroscopy)


$^1\text{H} - ^1\text{H}$

- HSQC (Heteronuclear Single Quantum Coherence)
- HMBC (Heteronuclear Multiple Bond Coherence)


$^1\text{H} - ^{13}\text{C}$

Why glycans are puzzling, why are they not obvious?


RNA




DNA



Protein



Glycan



Typical Applications of NMR:

- **Structural (chemical) elucidation**

Natural product chemistry

Organic chemistry: Analytical tool of choice for synthetic chemists.

- **Study of dynamic processes**

Reaction kinetics.

Study of equilibrium (chemical or structural).

- **Structural (three-dimensional) studies**

Proteins

DNA/RNA

Polysaccharides

Drug design

- **Structure Activity Relationships (SAR) by NMR**

- **Medicine - Magnetic Resonance Imaging (MRI)**

Structure Determination

- Each observable NMR resonance needs to be assigned or associated with the atom .
 - Molecular Formula
 - Functional groups
 - Carbon Connectivity
- Position of substitution on the carbon framework
 - Stereochemical properties.

NMR of carbohydrates:

- Sugar assignment (COSY, TOCSY, HSQC, HMBC)
- Anomeric configuration ($^3J_{\text{H,H}}$ coupling constants, NOE)
 - Ring size (^{13}C chemical shift, HMBC)
- Non carbohydrate substituents (Homo- and Heteronuclear NMR)
- Sugar sequence: Linkage analysis and glycosylation pattern (HMBC, NOE)

^1H and ^{13}C typical regions of carbohydrates:

The ^1H NMR Spectra can be roughly divided into the following regions:

Anomeric and Acylated Protons : 5.5-4.5 ppm.

Ring Protons : 4.5-3 ppm

Acetyl Groups, Methylene Protons: 3-2 ppm

Methyl Groups: 0.8-2.0 ppm

The ^{13}C NMR Spectra can be roughly divided into the following regions :

Anomeric Carbons Resonate Between 90-105 ppm

Ring Carbons Between 60-85 ppm

Nitrogen Bearing Carbons (In Amino Sugar) 50-60 ppm

Acetyl Groups 20-25ppm

Methylene Protons: 25-35 ppm


Methyl Groups: 15-20ppm

Sucrose

^1H

Anomeric region


Ring proton region




^{13}C




Anomeric region

Ring proton region







$$J = A - B \cos \phi + C \cos 2\phi$$




❖ Anomeric configuration

^1H NMR spectrum contains information on the configuration of glycosidic linkages




β -linkage: $J_{1,2} < 4$ Hz



β -linkage: $J_{1,2} > 6$ Hz


α -linkage: $J_{1,2} < 2$ Hz



α -linkage: $^1J_{\text{C}_1, \text{H}_1}$ 170-175 Hz
 β -linkage: $^1J_{\text{C}_1, \text{H}_1}$ 160-165 Hz


Sucrose

^1H




Ring proton region

$^3J_{\text{H1,H2}} 3.8 \text{ Hz} \implies \alpha$




Anatomy of a 2D NMR Experiment



2D NMR - The Interferogram


Interferogram



A 2D data set can be thought of as a series of 1D . Each 1D file is different from the next by a change in t_1 .

Fourier transformation of each 1D in the t_2 domain creates an interferogram.

2D plot of data





Two Dimensional NMR

- ✓ A 2D data set can be thought of as a series of 1D experiments collected with different timing.
- ✓ Fourier transformation of each 1D in the t_2 domain creates an interferogram.
- ✓ The t_1 domain is then Fourier transformed resulting in a 2D file with the frequency in each dimension.
- ✓ This 2D file will provide a map of all spin-to-spin correlations
- ✓ Each 2D experiment can provide either through bond (COSY type) or through space (NOESY type) correlation

COrrrelation SpectroscopY (COSY)

In a 2D COSY spectrum, **cross-peaks** will exist where there is spin-spin coupling between nuclei.




Cross peaks


Used to identify spins which are coupled to each other.

2D Experiments – COSY

The Power of 2D NMR:
Resolving Overlapping Signals




2 signals
overlapped



2 cross peaks
resolved

2D Experiments – COSY




Cross peaks due to geminal
and vicinal coupling



Sugar assignment


COSY of sucrose




Total Correlation Spectroscopy (TOCSY) experiment

TOCSY Experiment

In general, the TOCSY mixing time determines the number of bonds over which signal can be transferred, assuming that none of the coupling constants = 0





• Cross peaks generated between all members of a coupled spin network




TOCSY

- Cross peaks are generated between all members of a coupled spin network
- Coherence transfer period occurs during a multi-pulse spin-lock period;
- Length of spin-lock and J-coupling constants determine how far the spin coupling network will be probed



COSY and TOCSY – Sugar assignment




In Glucose, H1 and H2 protons are scalarly coupled, H1 and H3 are not.

In **COSY** spectra → H1 and H2 correlation observed ;

In **TOCSY** spectra → H1 and H3 observed


COSY and TOCSY – Sugar assignment

TOCSY of sucrose



HSQC: Heteronuclear Single-Quantum Correlation

Sucrose HSQC




The spectrum contains a peak for each unique proton attached to the heteronucleus being considered.

The 2D HSQC experiment permits to obtain a 2D heteronuclear chemical shift correlation map between directly-bonded ^1H and X-heteronuclei (an atomic nucleus other than a proton), often ^{13}C or ^{15}N .

HSQC: Heteronuclear Single-Quantum Correlation


^1H - ^{15}N HSQC spectrum of a fragment of the protein NleG₃-2.

Each peak in the spectrum represents a bonded N-H pair, with its two coordinates corresponding to the chemical shifts of each of the H and N atoms. Some of the peaks are labeled with the amino acid residue that gives that signal




HMBC (Heteronuclear Multiple Bond Correlation)


2D HMBC experiment correlates chemical shifts of two types of nuclei separated from each other with two or more chemical bonds.



HSQC and HMBC of Menthol



Long range *inter-residual* correlations in the HMBC spectrum




A₁ – B₃ long range correlations
(H₁ A – C₃ B and C₁ A – H₃ B)



A₁- B₃ linkage / A and B are 1-3 linked

Sugar Assignment, Ring size, Linkage analysis



HMBC of Sucrose



Nuclear Overhauser Effect (NOE) Spectroscopy


The 2D spectrum will have chemical shifts in f_1 and f_2 .

The cross peaks are for nuclei that are coupled dipolarly.





NOE contact: C is close in space to spin A

The NOE effect is the method for the elucidation of 3D structural features and stereochemistry



NOE and Distances

Isolated spin pair approximation (ISPA)




$$\frac{\sigma_{H1'-H4}}{\sigma_{H1'-H3'}} = \frac{r_{H1'-H4}^6}{r_{H1'-H3'}^6}$$

$$\eta_{ab} \propto r_{ab}^{-6}$$


$$\eta_{ac} \propto r_{ac}^{-6}$$

$$r_{ac} = r_{ab} * (\eta_{ab} / \eta_{ac})^{-1/6}$$




Intra-residue NOE contacts in monosaccharides: relative configuration of sugar residues

gluco, galacto configuration




α -linkage: H₁/H₂




β -linkage: H₁/H₃, H₁/H₅

manno configuration



α -linkage: no contact




β -linkage: H₁/H₂, H₁/H₃, H₁/H₅

Monosaccharide Sequence

❖ *Inter-residue NOE*


- ❖ Glycosylation shift (HSQC spectrum)
- ❖ Inter-residual long range correlation (HMBC spectrum)

Inter-residue NOE contacts in saccharides (Linkage analysis)




α -(1-3) linkage

β -(1-3) linkage



Sugar sequence - Linkage analysis


NOE in disaccharides may occur not only at the linkage protons but also at the neighbouring protons



...Saccharide conformation...


Sugar sequence - Linkage analysis

Sucrose **NOESY** and **TOCSY**



Application of various NMR techniques to carbohydrates


- HOMONUCLEAR (^1H - ^1H)
- HETERONUCLEAR (^1H - ^{13}C)




HOMONUCLEAR ^1H - ^1H
COSY
TOCSY
NOESY/ROESY
HETERONUCLEAR ^1H - ^{13}C
HMQC/HSQC
HMBC



A BACTERIUM'S ENEMY ISN'T YOUR FRIEND

a Strong immune response against WTA



b Weak immune response against WTA





Methicillin-resistant *Staphylococcus aureus* alters cell wall antigen glycosylation to subvert protective host defense



MOLECULAR INTERACTION by NMR

The ligand-based approach


- STD NMR
- TrNOESY
- WaterLOGSY
- Relaxation experiments
- Diffusion experiments

- Other nuclei: ^{19}F
- Paramagnetic tagging
- Other methods


The receptor based approach

- Isotope labelling
- Chemical shift perturbation mapping
- Paramagnetic tagging
- Other variations

Representation of protein-ligand interactions



Ligand observation



Two states equilibrium

L_{free} L_{bound} Molar fractions

FAST EXCHANGE

$$K_d = \frac{k_{\text{off}}}{k_{\text{on}}}$$

diffusion controlled

$$k_{\text{off}} = >10^2 \text{ (s}^{-1}\text{)}$$

$$k_{\text{on}} = >10^7 \text{ (s}^{-1}\text{M}^{-1}\text{)}$$

$$R_{\text{Lobs}} = L_f * R_{L_f} + L_b * R_{L_b}$$

$$\Delta R = L_b * (R_{L_b} - R_{L_f})$$

Experimental procedure: $L_0 \gg R_0$;

$$L_0/R_0 > 10 - 100 \dots \quad L_f \gg \gg L_b$$

Necessary condition: $|(R_{L_b} - R_{L_f})| \gg 0$

R_{L_b} Strong dependency on molecular size


NMR observable parameter R : **NOE; Diffusion; Line Shape**

Recapitulation

- K_{off} fast in the relaxation time scale, dissociation must occur before relaxation.
- K_{on} related to the efficacy of the interaction.
- Consider the molar fraction of ligand free and bound to the protein.
- Fast exchange in the chemical shift time scale
- **$L_0 \gg R_0$ means excess of free ligand but since we are in conditions in which the exchange is high (rate) the system is dominated by the bound state**
- K_{off} Dissociation before relaxation takes place; dissociation rate high since the molecule relax in the binding site

TRANSFERRED NOE

Information on the ligand bioactive conformation




- ❖ During the mixing time *inter* and *intra*-molecular NOE effects build up
- ❖ *Inter*-molecular tr-NOE effects are visible, intermolecular trNOEs are usually much larger than intramolecular effects


Important notes

- ❖ The mixing time must be short enough so that the contribution of the free ligand is negligible and long enough to allow visualization of the signal in the spectrum.
- ❖ The molar ratio of ligand to receptor. It should be emphasized that the trNOESY experiment works well for ligands that have K_D in the range 10^{-3} – 10^{-6} M / μM - mM range
- ❖ Small amount of purified receptor
- ❖ Routinely used to probe ligand-receptor interaction

TRANSFERRED NOE and MOLECULAR MOTION



The bioactive conformation: Transfer NOESY




🎯 Is There Any Binding?

🎯 Which Is The Ligand Bioactive Conformation?

Saturation Transfer Difference NMR Spectroscopy – STD NMR


IRRADIATION at the aromatic or aliphatic NMR regions

IS THERE ANY BINDING FOR ANY GIVEN COMPOUND?
WHICH IS THE BINDING EPITOPE?



At long irradiation times, the saturation is transferred to the bound ligand, first to the protons belonging to the ligand epitope, then to the rest of the ligand

Schematic representation of STD NMR method.



Key elements of protein-substrate binding


- H** close to receptor
- H** far from receptor
- H** furthest away from receptor

RECAPITULATION...

- **The saturation is transferred first to the protons belonging to the ligand epitope, then to the rest of the ligand**
- STD involves selective perturbation of protein-specific methyl proton resonances (0.5 p.p.m.–1 p.p.m.). This perturbation rapidly diffuses throughout the protein.
- This is usually done via a selective 180-degree pulse and results in a transfer of magnetization from the protein to any transiently bound ligands via the nuclear Overhauser effect (NOE).
- If the fragment binds to the target protein, the buildup of NOE that is transferred to the ligand results in enhanced signal corresponding to the resonances of that ligand in the STD spectrum.
- A number of factors affect the signal strength in an STD experiment, including protein size, duration of on-resonance irradiation, the frequency of irradiation, the dissociation constant of the ligand, the ligand/protein ratio and the field of the spectrometer.
- As such, it is useful to vary parameters such as ligand and protein concentration, the frequency of irradiation and the duration of irradiation

NMR Spectroscopy Techniques for Screening and Identifying Ligand Binding to Protein Receptors


β -Gal-Ome and RCA120 (Ricinus communis agglutinin), 50 μ M and 1.2 mM, 1:40



$$\text{STD effect: } (I_0 - I_{\text{sat}})/I_0$$

NMR Spectroscopy Techniques for Screening and Identifying Ligand Binding to Protein Receptors

β -Gal-Ome and RCA120 (Ricinus communis agglutinin), 50 μ M and 1.2 mM, 1:40




In order to determine the magnitude of the STD effects, the intensity of the signal in the STD NMR spectrum are compared with the signal intensities of a reference spectrum (off-resonance).

The STD signal with the highest intensity is set to 100% and the others are normalized to this signal.

Epitope Mapping with STD NMR Spectroscopy


Epitope mapping for sialyl LewisX bound to the the lectin *Aleuria aurantia* agglutinin (AAA)



STD TOCSY spectrum of sialyl LewisX in the presence of AAA (molar ratio 100:1): only the spin system of the l-fucose residue (F, yellow circle) is visible



TOCSY

STD-TOCSY



1D and 2D STD spectra shows unambiguously that only the fucose interact with the protein

Water-LOGSY NMR




The resonances of non-binding compounds appear with opposite sign and tend to be weaker than those of the interacting ligands.


Experimentally, the first step is the selective water excitation; during the mixing time (to be optimized based on the size of the complex) the water magnetization that has migrated to the protein is transferred to the ligand via direct or relay processes.

Carr Purcell Meiboom Gill (CPMG) experiments

CPMG, is a relaxation-time-edited NMR experiment that exploits differences in transverse relaxation time (T_2). *Proteins (and bound ligands) have a small T_2 while free ligands have a large T_2 . Thus monitoring T_2 , binding can be detected when the signal of the ligand decreases.*



In the CPMG experiment, we should see the signal decrease (or disappear) in the presence of protein and a binding fragment





In the presence of protein, the signal decreases, which is indicative of fragment binding. In the presence of both protein and a molecule known to bind the target (displacer), the signal is restored by ~30%. The protein target is *M. tuberculosis* CoaBC (10 μM). Fragment and displacer are present at 0.5 and 0.25 mM, respectively


Siglecs: sialic-acid binding immunoglobulin-like lectins

SIGLECS – SIALIC ACID AXIS


- Transmembrane proteins on innate immune cells
- Binding preferences for different modifications of Neu5Ac



H-CD22- sialoglycans studies




STD
Epitope
mapping



Bent conformation
 $\phi/\psi/\omega$ $-60^\circ/180^\circ/60^\circ$



CORCEMA-
st protocol

Bioactive
Conformation



Molecular
Dynamics



Docking
analysis




h-CD22- trisaccharide interaction: toward bioactive conformation

□ Omega angle


$$\omega = \text{O5-C5-C6-O6}$$



gt ω 60°





gg ω -60°



tg ω 180°


HSQC-TOCSY (blue-green)

HSQC-HECADE (red)



$$^2J_{(C6\ H5)} = -1.29 + 1.53\cos(\omega) - 3.68\sin(\omega)$$

Siglecs interaction with pathogens



- Cell signalling
- Cell adhesion
- Tumour sialic acids engagement
- Targets for cancer immunotherapy



- Uptake of sialylated pathogens



- Mechanisms to escape immune recognition



Dynamic changes of structures and composition of microbial cell envelope glycans, their **camouflage with host self-antigens (molecular mimicry)**, **their cross-talk with host immune Siglecs**, is central in GLYTUNES. The understanding of the role of sialic acid–Siglec interactions in the context of pathogen-associated immune suppression will pave the way to the **development of novel therapeutics glycomimetic to be used in clinical and diagnostic applications**

Questions?

